

THE OXIDATION OF ALCOHOLS AND ETHERS USING
CALCIUM HYPOCHLORITE [Ca(OCl)₂]

Stephen O. Nwaukwa and Philip M. Keehn

Tetrahedron Letters, Vol. 23, No. 1, pp 35-38, 1982

Summary: Calcium hypochlorite, a relatively stable, and easily stored and used solid hypochlorite oxidant, was found to oxidize secondary alcohols to ketones in excellent yields. Primary alcohols gave esters where both the acid and the alcohol portions of the ester were derived from the alcohol. Ethers were oxidized to esters though only in moderate yield.

While carrying out studies on the Grob-type cleavage of γ -hydroxy sulphones^{2,3} we attempted the preparation of the corresponding hypochlorite by the action of sodium hypochlorite on 2-tosylmethyl cyclohexanol⁴. Instead of the hypochlorite, an excellent yield of the corresponding ketone (2-tosylmethyl cyclohexanone) was obtained. Subsequent to this conversion, we independently found that this method was general for converting secondary alcohols to ketones by the use of sodium hypochlorite or commercially available Chlorox solutions. Though our conditions differed somewhat, the results were essentially the same as those recently reported by Stevens⁵.

The instability of sodium hypochlorite solutions however led us to consider other hypochlorite reagents which would be more stable and easier to handle. Calcium hypochlorite⁶ is a commercially available solid and is inexpensive. Since this reagent does not decompose significantly when stored without light in a desiccator⁷, carrying out oxidations by weighing the required amount of solid oxidant represented a more convenient method than using solutions which would frequently have to be titrated. We now wish to report our results concerning the use of Ca(OCl)₂ as an oxidant.

Oxidations of secondary alcohols with calcium hypochlorite proceeds smoothly, and in excellent yield at 0° in a solvent containing acetic acid. Our results are given in Table 1 for twelve compounds along with our initial

Table 1. The Oxidation of 2°-Alcohols Using Calcium and Sodium Hypochlorite.

Run	Substrate	Product	% Yield ^a		Reference
			Ca(OCl) ₂	NaOCl	
1	l-menthol	l-menthone	98	98	9a,b
2	borneol	camphor	98	99	9c
3	norborneol	norcamphor	92	--	10
4	cyclohexanol	cyclohexanone	91	98	9c
5	2-tosylmethyl- cyclohexanol ⁴	2-tosylmethyl- cyclohexanone ^c	98	98	--
6	3,5-dimethyl- cyclohexanol	3,5-dimethyl- cyclohexanone	93	--	10
7	5-cholesten-3-ol	4-cholesten-3-one	91	91	10 ^b
8	3-pentanol	3-pentanone	97	--	9d
9	3-pentanol	3-pentanone	--	87	9d
10	2-octanol	2-octanone	80	99	9d
11	diphenylcarbinol	benzophenone	98	--	9d
12	2-tosylmethyl-1- phenyl ethanol ⁴	α-tosylmethyl acetophenone ^d	--	98	--

a) Isolated yield

b) Methylene chloride was used as solvent instead of acetonitrile for solubility reasons.

c) Ir, CH₂Cl₂ (cm⁻¹) 1710, 1320, 1150; nmr, CDCl₃ (δ), 7.1-7.9 (m, 4H), 3.8-4.02 (dd, 2H) 2.5 (s, 3H), 1.5-2.1 (m, 9H); ms, m/e 266, 111.

d) Mp 130-131°; ir, CH₂Cl₂ (cm⁻¹) 1690, 1315, 1150; nmr, CDCl₃ (δ) 7.3-8 (m, 9H), 3.52 (s, 4H), 2.47 (s, 3H); ms, m/e 288.

results using sodium hypochlorite⁸. A general procedure is outlined for the oxidation of l-menthol to l-menthone. Thus l-menthol (3 g, 19 mmol) was dissolved in acetonitrile: acetic acid (3:2 25 ml) and added dropwise over a period of ten minutes to a cooled (0°C) and stirred solution of Ca(OCl)₂ (1.84 g, 12.7 mmol) in water (40 ml). Stirring was continued for 1 hr after which water (40 ml) was added. The solution was extracted with CH₂Cl₂ (4 x 30 ml) and the organic layers washed with 10% NaHCO₃ followed by an aqueous

Table 2. Oxidation of 1° Alcohols and Ethers Using Calcium and Sodium Hypochlorite.

Run	Substrate	Product	% Yield ^a		Reference
			Ca(OCl) ₂	NaOCl	
1	benzyl alcohol	benzaldehyde	98	98	10
2	1-pentanol	pentyl pentanoate	83	91	10
3	1-hexanol	hexyl hexanoate	98	98	9e
4	3-methyl butanol	3-methyl butyl isovalerate	76	87	10
5	ethyl alcohol	ethyl acetate	--	b	9b,10
6	ethyl ether	ethyl acetate	b	--	--
7	butyl ether	butyl butanoate	40	--	10
8	tetrahydrofuran	γ-butyrolactone	68	--	9b,10
9	tetrahydropyran	δ-valerolactone	56 ^c	--	9b,10

a) Isolated yield

b) Yield not calculated due to the volatility of the products but significant conversion was indicated by ir and nmr analysis.

c) Yield obtained by gc analysis.

wash. After drying with MgSO₄ and evaporating the CH₂Cl₂ the crude product was distilled affording *l*-menthone (2.89 g, 98%). The spectra (ir and nmr) were identical with those of authentic material^{9,10}.

Oxidation of primary alcohols under identical conditions gave an aldehyde only in the case of benzyl alcohol¹¹. Other primary alcohols gave esters as tabulated in Table 2. This table also includes our results on the oxidation of ethers to esters. Though the yields were not nearly as good as for the alcohols, the data is reported because of the unusual and potentially useful transformation¹². The ethers were oxidized under similar conditions as the alcohols except that the reactions were carried out at room temperature for from 4-16 hrs. Heating does not seem to increase the yield.

We are presently carrying out studies to improve the yields on the ether to ester transformation and to utilize the 1° alcohol oxidation for the preparation of lactones from α-ω diols.

REFERENCES

1. —
2. M. F. Semmelhack and J. C. Tomesch, J. Org. Chem., 42, 2657 (1977).
3. C. A. Grob and P. W. Schiess, Angew. Chem. Int. Ed. Engl., 6, 1 (1967).
4. The synthesis of this alcohol will be described in a forthcoming publication. The correct elemental analysis has been obtained.
5. R. V. Stevens, K. T. Chapman and N. H. Weller, J. Org. Chem., 45, 2030 (1980).
6. Obtained from Fisher Scientific Company, typical analysis 67% $\text{Ca}(\text{OCl})_2$.
7. Titrations were carried out over a period of two and one half months and there was no change in the concentration of the oxidant. Titrations were carried out as described in "A Textbook of Quantitative Inorganic Analysis", 3rd Edition by Arthur I. Vogel. Published by J. Wiley and Sons, New York, N.Y. (1961).
8. Both Chlorox (5.25% oxidant) and freshly prepared NaOCl were used without significant difference. NaOCl was prepared by bubbling chlorine into a solution of NaOH as described by V. Boido and O. E. Edwards, Can. J. Chem. 49, 2664 (1971).
9. a) C. J. Pouchert, The Aldrich Library of Infrared Spectra, Second Edition, published by Aldrich Chemical Co., Inc., Milwaukee, Wisconsin, 1975. b) C. J. Pouchert and John R. Campbell, The Aldrich Library of NMR Spectra Vol. II, published by Aldrich Chemical Co., Inc., Milwaukee, Wisconsin, 1974. c) Merck Index, 8th Edition, Merck and Co., Inc., N.J. 1968. d) ibid. 9th Edition, Merck, and Co., Inc., N.J., 1976. e) Chemical Abstract, 37, 5021.
10. Dictionary of Organic Compounds, 5th Edition, Oxford University Press, N.Y., 1969.
11. C. Y. Meyers, J. Org. Chem., 26, 1046 (1961).
12. a) L. M. Berkowitz and P. N. Rylander, J. Am. Chem. Soc., 80, 6682 (1958). b) Yoshiko Kamiya and S. Takemuro, Chem. Pharm. Bull., 21, 1401 (1973).
13. N. C. Demo and N. H. Potter, J. Am. Chem. Soc., 89, 3550 (1967).